

①  **Europäisches Patentamt**
European Patent Office
Office européen des brevets

⑪ Publication number:

0 145 294
A2

⑫

EUROPEAN PATENT APPLICATION

⑪ Application number: 84307788.8

⑤ Int. Cl.⁴: **C 07 D 249/08**
A 01 N 43/653

⑫ Date of filing: 09.11.84

⑬ Priority: 10.11.83 US 551732

⑭ Date of publication of application:
19.06.85 Bulletin 85/25

⑮ Designated Contracting States:
AT BE CH DE FR GB IT LI LU NL SE


⑰ Applicant: Rohm and Haas Company
Independence Mall West
Philadelphia, Pennsylvania 19105(US)

⑱ Inventor: Fujimoto, Ted Tsutomu
44 East Norton Drive
Churchville, Pa. 18966(US)

⑲ Representative: Angell, David Whilton et al,
Rohm and Haas Company Patent Department
Chesterfield House Bloomsbury Way
London WC1A 2TP(GB)

⑤ Substituted triazoles, processes for making them, their use as fungicides and fungicidal compositions containing them.

⑥ There are provided novel α -(halo-substituted phenyl)- α -alkyl- β -(1,2,4-triazol-1-yl)propanenitriles of Formula I below



wherein X is fluoro, chloro or bromo and R is (C₃-C₈)alkyl with the proviso that when R is branched chain (C₃-C₈)alkyl the branching does not occur at the α -carbon of the R substituent. There are also provided agronomically acceptable acid addition salts or agronomically acceptable metal salt complexes of a propanenitrile of Formula I. The above compounds are useful for combating phytopathogenic fungi.

EP 0 145 294 A2


0145294

SUBSTITUTED TRIAZOLES, PROCESSES FOR MAKING THEM, THEIR USE
AS FUNGICIDES AND FUNGICIDAL COMPOSITIONS CONTAINING THEM.

This invention concerns novel substituted triazoles,
processes for making them, their use as fungicides and
fungicidal compositions containing them.

U.S. Patent No. 4,366,165 and Mitsudera and Konishi
[J. Takeda Res. Lab., 41, 148-153 (1982)] disclose α -(halo-
substituted phenyl)- α -alkyl- β -(1,2,4-triazol-1-yl)propane-
nitriles and their use in combating phytopathogenic fungi.
However these references fail to point to the specific class
of compounds of the present invention which have a
particularly high degree of fungicidal activity against
certain fungi, notably those on wheat or barley.

The present invention provides α -(halo-substituted
phenyl)- α -alkyl- β -(1,2,4-triazol-1-yl)propanenitriles
characterized in that said propanenitriles are of Formula I
below



wherein X is fluoro, chloro or bromo and R is (C₃-C₈)alkyl
with the proviso that when R is branched chain (C₃-C₆)alkyl
the branching does not occur at the α -carbon of the R
substituent.


The invention also provides agronomically acceptable
acid addition salts and agronomically acceptable metal salt
complexes of compounds of Formula I.

The term "alkyl" used herein includes both branched
and straight chained alkyl groups but where a specific alkyl

0145294

group containing no prefix is referred to the n-alkyl group in question is meant, eg "butyl" means "n-butyl". Typical alkyl groups are propyl, n-butyl, iso-butyl, pentyl, neo-pentyl, iso-pentyl, hexyl, heptyl, octyl and iso-octyl.


An illustrative but preferred compound of Formula I has the formula:



This compound is named herein α -butyl- α -(4-chlorophenyl)- β -(1,2,4-triazol-1-yl)propanenitrile but it could alternatively be named α -butyl- α -(4-chlorophenyl)-1H-1,2,4-triazole-1-propanenitrile.

The acids which can be utilized in making the acid addition salts include, for example, hydrochloric, hydrobromic, nitric, sulfuric, phosphoric, hydroiodic, hydrofluoric, perchloric, p-toluenesulfonic, methanesulfonic, acetic, citric, tartaric, malic, maleic, oxalic, fumaric and phthalic acids.

Preferred metal salt complexes are of the Formula (II):



wherein X and R are as defined in Formula (I) above and M is a cation selected from Group IIA, IB, IIB, VIB, VIIB and VIII of the Periodic Table as shown in Handbook of Chemistry

and Physics and X is an anion selected so that the sum of the valence charges of the cation M and anion X equal zero. Typical cations are magnesium, manganese, copper, nickel, zinc, iron, cobalt, calcium, tin, cadmium, mercury, chromium, lead and barium. Typical anions are chloride, bromide, iodide, fluoride, sulfate, bisulfate, perchlorate, nitrate, nitrite, phosphate, carbonate, bicarbonate, acetate, citrate, oxalate, tartrate, malate, maleate, fumarate, p-toluenesulfonate, methanesulfonate, mono- or di-(C₁-C₄)alkyldithiocarbamate and (C₁-C₄)alkylene-bisdithiocarbamate.

Preferred compounds are those wherein in Formula I X is fluoro or chloro. It is also preferred that R is (C₃-C₆)alkyl. More preferred compounds are those where X is fluoro or chloro and R is a propyl, butyl, isobutyl or pentyl. The most preferred two compounds are those where X is chloro and R is n- or iso-butyl.

Typical compounds provided by the present invention include:

alpha-(4-chlorophenyl)-alpha-propyl-beta-(1,2,4-triazol-1-yl)propanenitrile
alpha-(4-fluorophenyl)-alpha-propyl-beta-(1,2,4-triazol-1-yl)propanenitrile
alpha-butyl-alpha-(4-chlorophenyl)-beta-(1,2,4-triazol-1-yl)propanenitrile
alpha-butyl-alpha-(4-bromophenyl)-beta-(1,2,4-triazol-1-yl)propanenitrile
alpha-butyl-alpha-(4-fluorophenyl)-beta-(1,2,4-triazol-1-yl)propanenitrile
alpha-(4-chlorophenyl)-alpha-iso-butyl-beta-(1,2,4-triazol-1-yl)propanenitrile
alpha-(4-chlorophenyl)-alpha-pentyl-beta-(1,2,4-triazol-1-yl)propanenitrile

0145294

alpha-(4-chlorophenyl)-alpha-isopentyl-beta-(1,2,4-
triazol-1-yl)propanenitrile

alpha-(4-fluorophenyl)-alpha-hexyl-beta-(1,2,4-
triazol-1-yl)propanenitrile

5 alpha-(4-fluorophenyl)-alpha-heptyl-beta-(1,2,4-
triazol-1-yl)propanenitrile


alpha-(4-chlorophenyl)-alpha-octyl-beta-(1,2,4-
triazol-1-yl)propanenitrile

10 and the agronomically acceptable acid addition salts and
metal salt complexes thereof.

The compounds provided by the present invention
possess curative, residual and preventive antifungal
properties against a broad spectrum of phytopathogenic
fungi. They additionally may act as systemic and/or contact
15 fungicides. Examples of such fungi include wheat and barley
powdery mildew (Erysiphe graminis), rice blast (Biricularia
oryzae), peanut early leaf spot (Cercospora arachidicola),
banana Sigatoka (Mycosphaerella fijiensis), wheat leaf rust
(Puccinia recondita), wheat stem rust (Puccinia graminis),
20 Septoriosis of wheat (Septoria spp.), barley net blotch
(Helminthosporium teres), grape powdery mildew (Uncinula
necator), grape black rot (Guignardia bidwellii), apple scab
(Venturia inaequalis), apple powdery mildew (Podosphaera
lencotricha), cucumber powdery mildew (Sphaerotheca
25 fuliginea), brown rot of stone fruits (Monilinia fructicola)
and rice sheath blight (Rhizoctonia solani).




The triazoles concerned in the present invention can
be prepared by conventional synthesis routes involving the
reaction of 1-H-1,2,4-triazole with one or more reagents
30 whereby in one or more steps there is introduced into the
1-position of the triazole a substituent of the formula:

0145294



For example, the triazoles may be prepared by nucleophilic displacement of the alkylated phenylacetone nitrile bromide (V) by a salt, preferably an alkali metal salt, of the triazole, generally about 1 to about 3 equivalents. This reaction can be run either neat or, preferably, in an appropriate solvent such as dimethylsulfoxide (DMSO), n-dimethyl-formamide, toluene or xylene at a temperature from about 0°C to about 150°C, preferably from about 25 to about 100°C. The bromide (V) is prepared by bromomethylation of the alkylated phenylacetone nitrile (IV) by methylenebromide (generally about 1.1 to about 2 equivalents) under basic conditions, e.g., sodium or potassium hydroxide, sodium or potassium hydride, potassium methoxide and potassium-t-butoxide (generally about 1.1 to about 2 equivalents) preferably with the use of a solvent such as DMSO, at a temperature from about 0 to about 150°C, preferably from about 25 to about 100°C. The alkylated phenylacetone nitriles (IV) can be prepared by phase transfer alkylation of the appropriately substituted benzylcyanides (III) with generally about 1 to about 2 equivalents of an alkyl halide (RX wherein R is as defined above and X is, for example, Cl, Br, I, tosylate or mesylate) in the presence of a strong base, e.g., 50% (w/w) sodium hydroxide or another metal alkoxide, and a catalyst, e.g., tetrabutylammonium bromide. Both the benzylcyanides and the alkyl halides can be readily prepared by techniques known from the literature. This synthesis scheme is shown below:

0145294



The acid addition salts of the triazoles can be prepared by standard techniques well-known in the art. For example, the triazole of formula (I) can be dissolved in an appropriate solvent such as diethyl ether, tetrahydrofuran, ethanol, methanol and the like or combinations thereof and treated with an equivalent or excess amount of a mineral or organic acid which may or may not be dissolved in an appropriate solvent, the mixture is then either cooled or evaporated to give the salt which can either be used as such or recrystallized from an appropriate solvent or combination of appropriate solvents.

The metal salt complexes of the triazoles of formula I can be prepared by adding dropwise, with stirring, a stoichiometric amount of a metal salt dissolved in an appropriate solvent or combination of solvents to a solution

0145294

of the triazole of formula (I) dissolved in a similarly appropriate solvent or combination of solvents. The reaction mixture is briefly stirred and the solvent is removed under reduced pressure to give the metal salt complex of the respective triazole.

The metal salt complexes can also be prepared by mixing stoichiometric or excess amounts of the metal salt and a triazole of formula (I) in the desired amount of solvent containing the appropriate adjuvants just prior to spraying the plants. Adjuvants that may be included in this "in situ" preparation may be detergents, emulsifiers, wetting agents, spreading agents, dispersing agents, stickers, adhesives, and the like which are used in agricultural applications.

Solvents that can be utilized in these procedures include any polar solvent, e.g., water, methanol, ethanol, isopropanol or ethylene glycol and any aprotic dipolar solvent, e.g., dimethylsulfoxide, acetonitrile, dimethylformamide, nitromethane or acetone.

The metal salt cations that can be used in these procedures can be selected from the group consisting of calcium, magnesium, manganese, copper, nickel, zinc, iron, cobalt, tin, cadmium, mercury, chromium, lead and barium. Any appropriate anion, e.g., chloride, bromide, iodide, sulfate, bisulfate, phosphate, nitrate, perchlorate, carbonate, bicarbonate, hydrosulfide, hydroxide, acetate, oxalate, malate or citrate may be utilized as the counterion in the metal salt.

The compounds of formula I possess an asymmetric carbon atom and thus exist as racemic mixtures. The d and l enantiomorphs in these racemic mixtures can be separated via standard techniques such as fractional crystallization with

0145294

d-tartaric acid, l-tartaric acid, l-quinic acid and the like followed by basification and extraction of the d or l enantiomorph free base. The invention, of course, includes all enantiomorphous forms.

5 The compounds of formula I, acid addition salts and metal salt complexes provided by the present invention are useful as agricultural fungicides and as such can be applied to various loci such as the seed, the soil (or other growth medium), the foliage or plant habitat. For such
10 purposes one or more of these compounds can be used in the technical or pure form as prepared or as formulations which contain the active ingredient in amounts up to 99% by weight but more usually up to 95%, eg 5-95%, by weight. The compounds are usually used in conjunction with an
15 agronomically acceptable carrier or diluent. Formulations can take the form of wettable powders, emulsifiable concentrates, dusts, granular formulations, aerosols, or flowable emulsion concentrates. In such formulations, the compounds are extended with a liquid or solid carrier or
20 diluent and, when desired, suitable surfactants are incorporated.

 It is usually desirable, particularly in the case of foliar spray formulations, to include adjuvants, such as wetting agents, spreading agents, dispersing agents,
25 stickers, adhesives and the like in accordance with agricultural practices. Such adjuvants commonly used in the art can be found in the John W. McCutcheon, Inc. publication "Detergents and Emulsifiers, Annual".

 In general, the compounds provided by this invention
30 can be dissolved in certain solvents such as acetone, methanol, ethanol, dimethylformamide or dimethyl sulfoxide and such solutions can be extended with water. The

0145294

concentrations of the solution can vary from about 1% to about 90% by weight with a preferred range being from about 5% to about 50% by weight.

5 For the preparation of emulsifiable concentrates, the compound can be dissolved in suitable organic solvents, or a mixture of solvents, together with an emulsifying agent which permits dispersion of the fungicide in water. The concentration of the active ingredient in emulsifiable concentrates is usually from about 10% to about 90% by weight and in flowable emulsion concentrates, this can be as high as about 75% by weight.

15 Wettable powders suitable for spraying, can be prepared by admixing the compound with a finely divided solid, such as clays, inorganic silicates and carbonates, and silicas and incorporating wetting agents, sticking agents, and/or dispersing agents in such mixtures. The concentration of active ingredients in such formulations is usually in the range of from about 20% to about 98%, preferably from about 40% to about 75%, by weight. A typical wettable powder is made by blending 50 parts of alpha-(4-chlorophenyl)-alpha-butyl-beta-(1,2,4-triazol-1-yl)propanenitrile, 45 parts of a synthetic precipitated hydrated silicon dioxide sold under the trademark Hi-Sil, and 5 parts of sodium lignosulfonate. In another preparation a kaolin type (barden) clay is used in place of the Hi-Sil in the above wettable powder, and in another such preparation 25% by weight of the Hi-Sil is replaced with a synthetic sodium silico aluminate sold under the trademark Zeolex 7.

30 Dusts are prepared by mixing the active triazole ingredient with finely divided inert solids which can be organic or inorganic in nature. Materials useful for this purpose include botanical flours, silicas, silicates,

0145294

carbonates and clays. One convenient method of preparing a dust is to dilute a wettable powder with a finely divided carrier. Dust concentrates containing from about 20% to about 80% by weight of the active ingredient are commonly made and are subsequently diluted to from about 1% to about 10% by weight use concentration.

0 The compounds provided by the invention can be applied as fungicidal sprays by methods commonly employed, such as conventional high-gallonge hydraulic sprays, low-gallonge sprays, air-blast spray, aerial sprays and dusts. The dilution and rate of application can be readily determined by one in the art depending upon the type of equipment used, the desired method, timing and frequency of applications, plants to be treated and diseases to be
5 controlled. Generally, however, the fungicidal compounds provided by the present invention will be applied in an amount of from about 0.1 to about 22 kg of active ingredient per hectare when applied foliarly or to the soil.

0 As a seed protectant, the amount of the compound coated on the seeds is usually from about 1.4 g to about 113 g ounces of active ingredient per hundred pounds of seed and preferably from 2.8 g to about 28 g per 45 kg of seed. As a soil or other growth medium fungicide the compounds can be incorporated in the growth medium or applied to its
5 surface usually at a rate of from about 0.06 to about 22 kg, preferably from about 0.02 to about 11 kg and more preferably from about 0.1 to about 3 kg of active ingredient per hectare. As a foliar fungicide, the compounds are usually applied to growing plants at a rate of from about
0 0.01 to about 11 kg, preferably from about 0.02 to about 6 and more preferably from about 0.03 to about 1 kg of active ingredient per hectare.

0145294

Fungicides which can be used in combination with the fungicides of this invention include all those listed in U.S. Patent No. 4,366,165.

The compounds of formula I and the acid addition salts and metal salt complexes thereof can be advantageously employed in various ways. Since these compounds possess broad spectrum fungicidal activity, they can be employed as fungicides in turf, fruit orchards, vegetable crops, cereal crops, golf course applications and the storage of cereal grain.

The following Examples 1-4 show the preparation of illustrative but preferred compounds of the invention. Example 5 gives fungicidal efficacy data.

Example 1: Preparation of alpha-butyl-alpha-(4-chlorophenyl)-beta-(1,2,4-triazol-1-yl)-propanenitrile

A) 1-bromo-2-cyano-2-(4-chlorophenyl)hexane

In a 300 ml three neck flask fitted with a mechanical stirrer, thermometer and addition funnel was added 20.8 g (0.1 mol) of α -(p-chlorophenyl)hexanenitrile, 34.8 g (0.2 mol) of methylene bromide and 50 ml of DMSO. To the reaction flask was added 24 ml of 50% (w/w) sodium hydroxide, dropwise over a 35 minute period. Upon completion of the reaction, it was quenched by adding 500 ml of water. The aqueous mixture was extracted three times with ether, then the combined ether extracts were washed three times with water and once with brine. The organic phase was dried over sodium sulfate, filtered and the solvent removed by rotary evaporation yielding 22.6 g of a yellow oil.

0145294

NMR: 60 MHz 0.8-2.0 δ (m, 9H), 4.0 δ (br s, 2H) and 7.4 δ (s, 2H).

B) Alpha-butyl-alpha-(4-chlorophenyl)-beta-(1,2,4-triazol-1-yl)propanenitrile

5 In a single neck 500 ml flask fitted with a magnetic stirrer and drying tube was added 26.6 g (0.09 mol) of 1-bromo-2-cyano-2-(4-chlorophenyl)hexane, followed by 19.0 g of potassium 1-H-1,2,4-triazole (0.18 mol) and 100 ml of DMSO. The reaction was stirred for about 48 hours at
10 room temperature and then stirred for about 24 hours at 70°C. The reaction was quenched by pouring it into 1500 ml of water. The aqueous mixture was extracted four times with ether and the combined ether extracts were washed twice with water and once with brine. The organic phase was dried
15 over sodium sulfate, concentrated and redissolved in a minimum amount of ether. Hexane was added until the solution was cloudy, and then the flask was placed in the freezer. The crystals which formed were filtered off and dried. The filtrate was concentrated to obtain additional
20 less pure material for a total of 13.0 g.

Elemental Analysis - Theoretical

(Found): C: 62.36(62.55, 62.38); H: 5.94(6.07, 6.00); N: 19.41(19.22, 19.35); Cl: 12.28(11.00, 11.90).

25 Example 2: Preparation of alpha-(4-chlorophenyl)-alpha-propyl-beta-(1,2,4-triazol-1-yl)propanenitrile

A) 2-(4-chlorophenyl)pentanenitrile

In a four neck one liter flask fitted with a reflux condenser, mechanical stirrer and addition funnel was placed 227.4 g (1.5 mol) of 4-chlorophenylacetonitrile, 235.6 g (3.0 mol) of 1-chloropropane and 4.8 g (0.015 mol) of tetrabutylammonium bromide. 300 g (3.75 mol) of 50%
30 (w/w) NaOH was added dropwise over 30 minutes. The reaction

0145294

mixture quickly exothermed to 50°C and the reaction flask was placed in a cold water bath until the reaction mixture cooled to 35°C. Thereafter, the reaction mixture was heated for an additional 4 hours at 50°C, then the reaction was quenched by adding 1500 ml of water. The mixture was extracted twice with ether and the organic phase was washed with water and then with 200 ml of 10% (w/w) hydrochloric acid. The ether was dried, filtered, concentrated and distilled. The distillate was collected, at 105°C at 0.5 mm, and two main fractions of 156.3 g (88% purity) and 88 g (60% purity) were obtained. The higher purity material was used in the bromomethylation of step B below.

NMR: 60 MHz ($d\text{-CHCl}_3$): 1.0-2.0 δ (m, 7H), 3.8-3.9 δ (t, 1H) and 7.5 δ (s, 4H).

B) 1-bromo-2-cyano-2-(4-chlorophenyl)-pentane

In a 2000 ml four neck flask fitted with a reflux condenser, mechanical stirrer, thermometer and addition funnel was placed 305.0 g (1.57 mol) of 2-(4-chlorophenyl)pentanenitrile, 409.4 g (2.36 mol) of dibromomethane and 400 ml of DMSO. 50% (w/w) sodium hydroxide (251.2 g, 3.00 mol) was added dropwise over 2 hours with the reaction temperature rising to 50°C. A water bath was used to maintain the reaction at about 40°C and after about 4 hours, the reaction was quenched by adding 100 ml of water. After sitting at room temperature for about 16 hours, the mixture was diluted with 1000 ml of ice water, extracted with ether, the combined ether extracts were washed twice with water and then with 10% (w/w) hydrochloric acid and dried over sodium sulfate. The ether was removed by rotary evaporation and the remaining DMSO removed under high vacuum to yield 361.2 g of the bromide derivative.

0145294

NMR: 60 MHz ($d\text{-CHCl}_3$) 1.0-2.2 δ (m, 7H),
4.0 δ (br s, 2H) and 7.5 δ (s, 4H).

C) alpha-propyl-alpha-(4-chloro)-beta-(1,2,4-triazol-1-yl)propanenitrile

5 In a single neck one liter flask was placed
361.2 g (1.26 mol) of 1-bromo-2-cyano-2-(4-chlorophenyl)-
pentane and 300 ml of DMSO. The potassium salt of
10 1-H-1,2,4-triazole (270.0 g, 2.52 mol) was added and the
flask was placed on a rotary evaporator for 1 hour at 75°C
to dissolve the triazole salt. The flask was then heated at
90°C for 4 hours, followed by stirring at room temperature
overnight (about 16 hours). The reaction was found to be
incomplete and reheated to 120°C for about 12 hours followed
by stirring it at 90°C for about 36 hours. Upon completion
15 of the reaction, the reaction mixture was cooled to room
temperature and diluted with 1500 ml of ice water. The
water was extracted with 250 ml of ether; however, since the
product crystallized in the ether layer, this layer was then
diluted with 1500 ml of ethyl acetate and extracted. The
20 water layer was additionally extracted twice with ethyl
acetate and the combined organic extracts were separated
into two batches. Each was extracted with water
(6 x 150 ml) and once with brine. The dark brown organic
phase was dried over sodium sulfate and then filtered. The
25 solution was treated then with activated charcoal and
filtered and stripped to give a crystalline cake which was
recrystallized from ethyl acetate: ether. The product was
kept in the freezer for 16 hours and then filtered and
washed with hexane. The filtrate was concentrated, hexane
30 added and then triturated to yield an additional solid which
was combined with the first crop to give 166.8 g of product.

0145294

M.P.: 108-110°C

Elemental Analysis - Theoretical

(Found): C: 61.20(61.16); H: 5.50(5.56); N:
20.39(20.26); Cl: 12.91(12.66).

Example 3: Preparation of alpha-butyl alpha-(4-fluorophenyl)-beta-(1,2,4-triazol-1-yl)-propanenitrile

A) 2-(4-fluorophenyl)hexanenitrile

The phase transfer procedure was employed as with 2-(4-chlorophenyl)pentanenitrile, with 400 g (2.96 mol) of p-fluorophenylacetonitrile, 548 g (5.92 mol) of 1-chlorobutane, 592 g (7.4 mol) of 50% (w/w) sodium hydroxide and 9.07 g (0.3 mol) of tetrabutylammonium bromide. The base was added over 2.5 hours with the reaction temperature rising to 35°C. The reaction was then stirred for about 12 hours at 45°C. Upon completion, the reaction was quenched with water and the mixture was extracted and washed as in Example 1 and then distilled to yield 358.8 g of 91% pure product.

NMR data: 60 MHz ($d\text{-CHCl}_3$): 0.8-2.0 δ (m, 9H), 3.7-3.8 δ (t, 1H) and 7.4-7.5 δ (br s, 4H).

B) 1-bromo-2-cyano-2-(4-fluorophenyl)hexane

The bromomethylation procedure of Example 1 was employed using 358.1 g (1.875 mol) of 50% (w/w) sodium hydroxide, 488.9 g (2.81 mol) of dibromomethane and 400 ml of DMSO. Sodium hydroxide (50%, w/w) was added dropwise over about 2 hours with the reaction exotherming to 90°C. The reaction mixture was cooled to 50°C and the mixture was stirred for 11 hours until the reaction was about 90% complete. The reaction was quenched and worked up in the manner described in Example 1A to yield 379.5 g of a 88% pure product. The product was used directly to prepare the triazole adduct as described in step C below.

0145294

NMR: 0.8-2.0 δ (m, 9H), 4.0 δ (br s, 2H),
and 7.4 δ (s, 4H).

C) Alpha-butyl-alpha-(4-fluorophenyl)-beta-
(1,2,4-triazol-1-yl)propanenitrile

5 In a 2 liter four neck flask was placed
379.5 g (1.33 mol) of 1-bromo-2-cyano-2-(4-fluorophenyl)-
hexane and 300 ml of DMSO to which 156 g (1.46 mol) of
potassium 1-H-1,2,4-triazole was added. The reaction mixture
was stirred overnight at 90°C. The product was worked up in
10 the manner described in Example 1B to yield 340.5 g of a
semisolid, 83% pure.

NMR data: 60 MHz (d -CHCl₃): 1.0-2.2 δ
(m, 9H), 4.8 δ (s, 2H), 7.0-7.8 δ (m, 4H), 8.05 δ (s, 1H) and
8.10 δ (s, 1H).

15 Example 4: Preparation of alpha-(4-fluorophenyl)-alpha-
propyl-beta-(1,2,4-triazol-1-yl)-
propanenitrile

A) 2-(4-fluorophenyl)pentanenitrile

20 To 50.0 g (0.37 mol) of 4-fluorophenyl-
acetonitrile, 0.6 g of tetraethyl ammonium bromide, 50 g
(0.407 mol) of 1-bromopropane and 50 ml of DMSO was added
35.5 g (0.44 mol) of 50% (w/w) sodium hydroxide at room
temperature. After the addition was complete, the reaction
was stirred at 50-60°C for 3 hours. The reaction mixture
25 was extracted with ether, the organic phase washed with
water and then 10% (w/w) hydrochloric acid. The ether was
dried, filtered and distilled to give 23 g of the product
(b.p. 105-110°C at 5 mm).

30 NMR: 60 MHz (d -CHCl₃) 0.9-2.2 δ (m, 7H),
3.8-4.0 δ (t, 1H), 7.0-7.6 δ (m, 4H).

B) 1-bromo-2-cyano-2-(4-fluorophenyl)pentane
The bromomethylation of the previous

0145294

examples was employed using 23.0 g (0.129 mol) of 2-(4-fluorophenyl)pentanenitrile and 22.5 g (0.129 mol) of dibromoethane in 50 ml of DMSO. To the reaction mixture was added 11.95 g (0.15 mol) of 50% (w/w) sodium hydroxide while stirring at room temperature. The reaction was heated to 60°C for 3 hours, cooled to room temperature and worked up in the manner described in the previous examples, e.g., Example 3B. There was obtained 20.5 g of the bromide derivative which was used directly below in step C.

b.p. 105-110°C at 5 mm

NMR: 0.7-2.2 δ (m, 7H), 3.9 δ (q, 2H), and 7.0-7.8 δ (m, 4H).

C) Alpha-(4-fluorophenyl)-alpha-propyl-beta-(1,2,4-triazol-1-yl)propanenitrile

To a solution of 8.1 g (0.76 mol) of potassium 1-H-1,2,4-triazole and 30 ml of DMSO was added 15 g (0.58 mol) of 1-bromo-2-cyano-2-(4-fluorophenyl)pentane while maintaining the temperature at 50°C. While stirring, the reaction mixture was heated further to 80°C for 1 hour and then heated at 100°C for 24 hours. The product was worked up as described in Example 3C and after concentration and trituration, 1.3 g of a white solid, m.p. 7-76°C, was obtained.

Elemental Analysis - Theoretical

(Found): C: 65.09 (64.80); H: 5.86 (5.95); N: 21.69 (20.77); F: 7.36 (7.02).

NMR: 90 MHz ($d\text{-CHCl}_3$) 0.9-1.0 δ (t, 3H) 1.2-1.8 δ (m, 2H), 1.9-2.2 δ (t, 2H), 3.7-3.9 δ (q, 2H), 7.0-7.5 δ (m, 4H) and 7.9 δ (br s, 2H).

Following the procedures above-described there may also be prepared compounds of Formula I (and the acid addition salts and metal salt complexes), wherein (a) X is bromo and R is each and every one of propyl, butyl,

iso-butyl, pentyl, iso-pentyl, hexyl, iso-hexyl, heptyl, 2-heptyl, octyl or heptyl substituted with methyl in the 3-, 4- or 5-positions, (b) X is chloro and R is each and every one of iso-butyl, pentyl, iso-pentyl, hexyl, iso-hexyl, heptyl, 2-heptyl, octyl and heptyl substituted with methyl in the 3-, 4- or 5- positions and (c) X is fluoro and R is each and every one of the alkyl substituents listed in (b) immediately above.

Example 5

The compounds of Examples 1-4 were tested for their antifungal activity against wheat leaf rust (WLR), wheat powdery mildew (WPM) and barley spot blotch (BSB). The test procedures were as follows:

A) Wheat Leaf Rust (Puccinia recondita):

Pennoll wheat seedlings about 7 days old were sprayed to runoff with a solution of the test compound comprising 75 parts per million of the test compound suspended in a 2:1:1 by volume mixture of water, acetone and methanol. After drying, the plants were inoculated with a uredial suspension of P. recondita (20,000 uredial/ml). The plants were incubated at 22°C in a mist for 24 hours. After seven days of further incubation on a greenhouse bench, the percent disease control was determined by counting uredial pustules and comparing it to the number of pustules on control plants.

B) Wheat Powdery Mildew (Erysiphe Graminis

f. sp. tritici): Pennoll wheat seedlings (7-14 days old) were trimmed to provide a uniform plant height and to facilitate uniform inoculation. Twenty-four hours after their trimming, they were sprayed to runoff with a test compound comprising 5 parts per million of the test compound suspended in a 2:1:1 mixture of water, acetone and methanol. Then mildew spores of wheat powdery mildew cultured on wheat

0145294

seedlings in a controlled temperature room were shaken from the cultured plants onto the Pennoll wheat seedlings. The inoculated seedlings were kept in the controlled temperature room and subirrigated. The percent disease control was rated 8-10 days after inoculation.

C) Barley Spot Blotch (Helminthosporium sativum): Pennrad barley seedlings about 7 days old were sprayed to runoff with a test compound comprising 10 parts per million of the test compound suspended in a 2:1:1 by volume mixture of water, acetone and methanol. After drying, the plants were inoculated with a conidial suspension of H. sativum (20,000 conidia/ml). The plants were incubated at 22°C in a mist for 24 hours. After five days of further incubation on a greenhouse bench, the percent disease control was measured by counting lesions and comparing it to the number of lesions on the control plants.


The results of the tests are presented below in Table 1. The disease control ratings utilized in Table 1 are as follows: A = 97-100% disease control, B = 90-96% disease control, C = 70-89% disease control, D = 50-69% disease control and E = 49% or less disease control.

TABLE 1

<u>Compound</u>	<u>Disease Control Rating</u>		
	<u>WLR</u>	<u>WPM</u>	<u>BSB</u>
1	A	A	B
2	A	A	A
3	A	B	B
4	A	A	C

CLAIMS:

1. An α -(halo-substituted phenyl)- α -alkyl- β -(1,2,4-triazol-1-yl)propanenitrile, characterized in that said propanenitrile is of Formula I below



- 5 wherein X is fluoro, chloro or bromo and R is (C₃-C₈)alkyl with the proviso that when R is branched chain (C₃-C₆)alkyl the branching does not occur at the α -carbon of the R substituent; or an agronomically acceptable acid addition salt or an agronomically acceptable metal salt complex of a propanenitrile of Formula I.
- 10 2. A compound as claimed in Claim 1, wherein X is fluoro or chloro.
3. A compound as claimed in Claim 1 or 2, wherein R is (C₃-C₆)alkyl.
- 15 4. A compound as claimed in Claim 3, wherein R is straight chained alkyl.
5. A compound as claimed in Claims 2 and 3, wherein R is n-propyl, n-butyl, iso-butyl or n-pentyl.
- 20 6. A compound as claimed in Claim 3, wherein X is chloro and R is n-propyl or X is fluoro in which case R is n-propyl or n-butyl.
7. A compound as claimed in Claim 3, wherein X is chloro and R is n-butyl.

0145294

8. A fungicidal composition containing as active ingredient at least one compound as claimed in any preceding claim and an agronomically acceptable carrier or diluent for the active ingredient.

5 9. A method for combating phytopathogenic fungi which comprises applying to a plant, to plant seed, to a plant habitat or to growth medium a fungicidally effective amount of a compound as claimed in any one of Claims 1-7.

10 10. A method as claimed in Claim 9 as applied to combating fungi on wheat.

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 84307788.8

(51) Int. Cl.⁴: **C 07 D 249/08**
A 01 N 43/653

(22) Date of filing: 09.11.84

(30) Priority: 10.11.83 US 551732

(43) Date of publication of application:
19.06.85 Bulletin 85/25

(88) Date of deferred publication of search report: 26.02.86

(84) Designated Contracting States:
AT BE CH DE FR GB IT LI LU NL SE


(71) Applicant: Rohm and Haas Company
Independence Mall West
Philadelphia, Pennsylvania 19105(US)

(72) Inventor: Fujimoto, Ted Tsutomu
44 East Norton Drive
Churchville, Pa. 18966(US)

(74) Representative: Angell, David Whilton et al,
Rohm and Haas Company Patent Department
Chesterfield House Bloomsbury Way
London WC1A 2TP(GB)

(54) Substituted triazoles, processes for making them, their use as fungicides and fungicidal compositions containing them.

(57) There are provided novel α -(halo-substituted phenyl)- α -alkyl- β -(1,2,4-triazol-1-yl)propanenitriles of Formula I below



wherein X is fluoro, chloro or bromo and R is (C₃-C₈)alkyl with the proviso that when R is branched chain (C₃-C₈)alkyl the branching does not occur at the α -carbon of the R substituent. There are also provided agronomically acceptable acid addition salts or agronomically acceptable metal salt complexes of a propanenitrile of Formula I. The above compounds are useful for combating phytopathogenic fungi.



European Patent
Office

EUROPEAN SEARCH REPORT

0145294
Application number

EP 84 30 7786

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
X,Y	FR-A-2 391 200 (ROHM AND HAAS) * Page 6, lines 7-33; claims * & US - A - 4 366165 (Cat. D)	1-10	C 07 D 249/08 A 01 N 43/653
X,Y P	--- CHEMICAL ABSTRACTS, vol. 101, no. 23, 3rd December 1984, page 652, no. 211152t, Columbus, Ohio, US; & JP - A - 59 104 367 (TAKEDA CHEMICAL INDUSTRIES LTD) 16-06-1984 * Abstract *	1-10	
X,Y	--- EP-A-0 052 424 (ICI) * Claims 1-4,11,12 *	1-10	
Y,D	--- CHEMICAL ABSTRACTS, vol. 98, no. 25, 20th June 1983, page 556, no. 215541m, Columbus, Ohio, US; H. MITSUDERA et al.: "Synthesis and fungicidal activity of 1,2,4-triazole derivatives", & TAKEDA KENKYUSHOHO 1982, 41(3/4), 148-53 * Abstract *	1-10	TECHNICAL FIELDS SEARCHED (Int. Cl.4) C 07 D 249/00
Y	--- FR-A-2 300 081 (ROHM AND HAAS) * Pages 71-74; pages 80-82; page 90; claims * -----	1-10	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 26-11-1985	Examiner THEUNS H.G.
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			